Case Report

DOI:10.22374/jeleu.v2i1.23

ECTOPIC DECIDUAL REACTION IN BLADDER

Blessing Dhliwayo¹ and Ananda Kumar Dhanasekaran²

¹Royal Preston Hospital, Preston, Lancashire, United Kingdom

²City Hospital, Dudley Road, Birmingham, United Kingdom.

Corresponding Author Ananda Kumar Dhanasekaran: managestones@gmail.com

Submitted: October 23, 2018. Accepted: November 19, 2018. Published: January 24, 2019.

CASE PRESENTATION

A 32-year-old teacher, 12-weeks' pregnant presented with visible hematuria to midwife-led clinics. Ultrasound scans showed a single viable fetus and irregular

lesion at the left side of the bladder. It measured $28 \times 26 \times 34$ mm with presence of internal Doppler vascularity raising high suspicion of urological cancer. Ultrasound Doppler longitudinal with cross section images are shown in Figures 1 and 2. Both the kidneys

FIG. 1 Ultrasound Doppler.

FIG. 2 Ultrasound Doppler.

were normal. She was referred to urology. Her flexible cystoscopy showed irregular lesion covering left ureteric orifice coinciding with the ultrasound findings (Figure 3). Urine microscopy has no signs of infection. Urine cytology has no malignant cells. Findings were discussed in Urology Multidisciplinary Meeting. Considering the nature of the lesion transurethral resection of bladder lesion was planned.

DIAGNOSTIC CONUNDRUM

Informed consent obtained. Antenatal team involved to help in pregnancy related support. Under spinal anesthesia transurethral resection of bladder lesion done. Both the ureteric orifices were normal. She had uneventful recovery and sent home in 24 hours. Bladder fragments histology showed normal urothelium. Lamina propria contains nodular aggregates of decidual tissue with cells having abundant eosinophilic

cytoplasm, vesicular nuclei and inconspicuous nucleoli. Some degenerative changes are also seen in some of the decidual cells. No endometrial tissue or chorionic villi are seen. Diagnosis of ectopic decidual reaction in bladder was made. Histology in 4× shown in Figure 4 and 20× shown in Figure 5. She was updated on the good news of benign histology. She has completed her pregnancy uneventfully.

LITERATURE REVIEW AND DISCUSSION

The decidual reaction is the set of changes in the endometrium of the uterus that prepare it for implantation of an embryo due to progesterone. These changes are observed in each menstrual cycle of female and enhanced after implantation. These changes include swelling of stromal cells due to accumulation of glycogen and other nutrients. Ectopic decidual reaction is well explained in literature in intraperitoneal locations

FIG. 3 Flexible cystoscopy showed irregular lesion covering left ureteric orifice.

FIG. 4 Histology in 4×.

and genital tracts. But ectopic decidual reaction in bladder mimicking bladder growth is extremely rare. It is a difficult differential diagnosis because of its extreme rarity.

The first report of decidual transformation outside of the uterine mucosa was made by Walker in 1887,

FIG. 5 Histology in 20×.

who observed it in association with an extra-uterine pregnancy.¹ Taussig² coined the term ectopic decidua to describe the phenomenon in its extra-mucosal locations, and ectopic decidua has been noted in association with pregnancy in such diverse locations as the endocervical stroma, the sub-serosa of the corpus uteri and its ligamentous attachments, the endosalpinx, the serosal surfaces of the fallopian tubes, ovaries, omentum, vermiform appendix, small and large intestine, urinary bladder, and portions of the mesenteries. Meyer³ stressed the role of healed, low-grade, and pre-existing inflammation, especially that associated with the formation of adhesions, in the sensitization of the tissues for the decidual response. Ectopic decidua in pregnancy has been observed in the umbilicus, in laparotomy and hysterotomy scars, and in association with adenomyosis and endometriosis. Weller⁴ has pointed out that the regional distribution of the ectopic decidual response corresponds with the distribution of endometriosis. William B. Ober et al.⁵ published case series of 16 patients in 1957.

Only one case report was published in 2009 for bladder submucosal ectopic decidual reaction.⁶ In this case report we present florid papillomatous ectopic decidual reaction making difficult to differentiate from bladder transitional cell carcinoma. We hope this article will create awareness of this entity and help to treat similar patients in future. Since this bladder lesion resembles bladder cancer knowledge about ectopic

decidual reaction in bladder will avoid unnecessary apprehension and interventions. The learning point in this case report is to be aware of rare differential diagnosis of ectopic decidual reaction in pregnancy in pregnant patients presenting with bladder lesions.

REFERENCES

1. Walker A. Der Bau der Eihdute bei Graviditas abdominalis. *Virchows Arch. f. path. Anat* 1887;107:72–99.
2. Taussig FJ. Ectopic decidua formation. *Surg Gynec & Obst* 1906;2:292–303.
3. Meyer R. Die Entzündung als Entstehungsursache ectopischer Decidua oder Paradecidua. *Ztschr. f. Geburtsh. u. Gynaik.* 1913;74:250–77.
4. Weller CV. The ectopic decidual reaction and its significance in endometriosis. *Am J Pathol* 1935;11:287–290.
5. Ober WB, Grady HG, Schoenbucher AK. Ectopic ovarian decidua without pregnancy. *Am J Pathol* 1957;33:199–217.
6. Szopiński TR, Sudół-Szopinska I, Dzik T, Borówka A. Ectopic decidual reaction in the urinary bladder presenting as a vesical tumor. *Urology* 2009;74:1232–33.